


FAST 2024 FOOD ALLERGY TREATMENT SURVEY

Spring, 2024


Food allergens used for OIT

Answered: 106 Skipped: 13




The youngest age that you routinely accept for OIT.

Answered: 106 Skipped: 13




OIT starting dose

Answered: 106 Skipped: 13




Omalizumab during OIT

Answered: 106 Skipped: 13




Most Commonly Used Dose escalation frequency

Answered: 106 Skipped: 13




After Hours Management

Answered: 106 Skipped: 13




Q10: We treat peanut allergy using OIT

Answered: 107 Skipped: 12




We start with Palforzia

Answered: 96 Skipped: 23




Typical Day One Peanut Starting Dose in mg of peanut protein

Answered: 98 Skipped: 21




Day One Peanut Planned Final Dose in mg of protein

Answered: 96 Skipped: 23




Peanut target dose in patients <4 years of age in mg of peanut protein?

Answered: 94 Skipped: 25




Peanut target dose in patients >4 years of age in mg of peanut protein?

Answered: 96 Skipped: 23




Peanut Maintenance Dose patients <4 years of age

Answered: 93 Skipped: 26




Peanut Maintenance Dose patients >4 years of age

Answered: 95 Skipped: 24




Maintenance Product for Patients who used Palforzia for dose escalation

Answered: 94 Skipped: 25




We treat tree nut or seed allergy with OIT

Answered: 97 Skipped: 22




Tree nut and seed procedures

Answered: 85 Skipped: 34




Day One Tree Nut/Seed Starting Dose in mg of nut protein

Answered: 85 Skipped: 34




Day One Tree nut/Seed final dose in mg of protein

Answered: 85 Skipped: 34




Maintenance Dose for Tree nuts/Seeds for patients <4 years of age (mg of nut/seed protein)

Answered: 84 Skipped: 35




Maintenance Dose for Tree nuts/Seeds for patients >4 years of age (mg of nut/seed protein)

Answered: 85 Skipped: 34




Q25: We have treated milk allergy with OIT

Answered: 95 Skipped: 24




Day One Milk Starting Dose in mg of protein

Answered: 70 Skipped: 49




Day One Milk Final Dose in mg of protein

Answered: 69 Skipped: 50




Do you commonly permit baked or partially cooked milk to be used for milk up dosing?

Answered: 70 Skipped: 49




Milk Maintenance Dose for patients <4 years of age

Answered: 69 Skipped: 50




Milk Maintenance Dose for patients >4 years of age

Answered: 71 Skipped: 48




Q31: We have treated egg allergy with OIT

Answered: 95 Skipped: 24




Day One Egg Starting Dose in mg of protein

Answered: 68 Skipped: 51




Day One Egg Final Dose in mg of protein

Answered: 69 Skipped: 50




Do you commonly permit baked or partially cooked egg to be used for updosing?

Answered: 68 Skipped: 51




Maintenance Dose for Egg for patients <4 years of age

Answered: 67 Skipped: 52




Maintenance Dose for Egg for patients >4 years of age

Answered: 70 Skipped: 49




Multi-food OIT

Answered: 70 Skipped: 49




The minimum time taking the post-escalation dose before consideration of less frequent than daily dosing.

Answered: 92 Skipped: 27




Target and maintenance dose variation

Answered: 93 Skipped: 26




Modification of Maintenance Dose frequency

Answered: 93 Skipped: 26




The lowest maintenance dosing frequency prior to a sustained unresponsiveness challenge

Answered: 92 Skipped: 27




Post-dose maintenance activity limitation

Answered: 93 Skipped: 26




For maintenance patients who are avoiding their food allergen, we modify the 2 hour activity limitation after OIT dosing

Answered: 92 Skipped: 27




For maintenance patients who are "free eating" their allergenic food, we modify the 2 hour activity limitation after OIT dosing as follows:

Answered: 88 Skipped: 31




For maintenance patients who are "free eating" their allergenic food, we modify the 2 hour activity limitation after consuming the food as part of the regular diet:

Answered: 89 Skipped: 30




Sustained Unresponsiveness (SU) challenges

Answered: 91 Skipped: 28




Number of Sustained Unresponsiveness Challenges Performed

Answered: 78 Skipped: 41




Criteria for sustained unresponsiveness challenge

Answered: 64 Skipped: 55




Duration of avoidance prior to Sustained Unresponsiveness challenges

Answered: 60 Skipped: 59




Final dose of OIT food used for sustained unresponsiveness challenge

Answered: 61 Skipped: 58




Sustained unresponsive challenge pass rate

Answered: 47 Skipped: 72




Frequency of OIT food exposure after passing an SU challenge (avoiding the food)

Answered: 18 Skipped: 101




Frequency of OIT dosing after passing an SU challenge if the food is incorporated into the diet

Answered: 17 Skipped: 102




Epinephrine recommendation after passing a sustained unresponsiveness challenge

Answered: 18 Skipped: 101




Q55: Food SLIT

Answered: 94 Skipped: 25




Q57: SLIT starting age - the youngest age that you routinely accept for SLIT. If you do not offer SLIT, skip to the end of the survey.

Answered: 46 Skipped: 73




Q58: List all of the food allergens that you have used for SLIT. If you do not offer SLIT, skip to the next question.

Answered: 42 Skipped: 77




Q59: In our practice, indications for food SLIT (now or when we start in the next six months) include:

Answered: 48 Skipped: 71




Q60: We offer SLIT for the following food (check all that apply):

Answered: 45 Skipped: 74




Q61: We use

Answered: 42 Skipped: 77




Q62: If you use commercial allergen extracts, what dilution do you use for your SLIT starting dose

Answered: 25 Skipped: 94




Q63: If you use commercial allergen extracts, what dilution do you use for your SLIT maintenance dose

Answered: 21 Skipped: 98




Q64: If you use retail food for SLIT, what is your starting dose in mg of protein

Answered: 35 Skipped: 84




Q65: If you use retail food for SLIT, what is your maintenance dose in mg of protein

Answered: 35 Skipped: 84



Q66: Multi-food SLIT

Answered: 40 Skipped: 79



Q67: What is the exercise limitation post-SLIT dose

Answered: 41 Skipped: 78

